




Findings to date: *Enterococcus* spp. concentrations were an order of magnitude higher at the Puakō anchialine ponds compared to the other watershed locations (average  $\pm$ SE: 8165  $\pm$  2935 MPN/100



**Figure 1.** Spatial distribution of sewage indicators in the Puakō watershed, including shoreline stations at adjacent resorts north (Mauna Kea, Hapuna Prince) and south (Fairmont Orchid, Mauna Lani) of the community on Hawai‘i Island. A) *Enterococcus* spp., B) *Clostridium perfringens*, C) Human-associated *Bacteroides*, and D)  $\delta^{15}\text{N}$ . Puakō watershed samples were collected from June – August 2016; resort samples were collected from January – March 2017. Sample analysis for *Bacteroides* at the resorts is pending.

**Table 1.** Average  $\pm$  SE of *Enterococcus* spp., *Clostridium perfringens*, human-associated *Bacteroides* (markers HF183 and BacHum) and  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  and  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  in water samples for high elevation wells (Waikoloa Village), mid elevation wells (Mauna Lani), low elevation wells (Puakō wells), resort shoreline stations (Mauna Kea, Hapuna Prince, Fairmont Orchid, and Mauna Lani), Puakō shoreline stations (low and high salinity stations), and Puakō ponds (anchialine ponds). Number of stations sampled within each watershed category varied between 3-16 (station n) and the total number of observations per location per parameter are footnoted below. Water samples collected at the resorts for human-associated *Bacteroides* analyses have not yet been completed (n/a).

Watershed Location	Station (n)	<i>Enterococcus</i> spp. (MPN/100mL)	<i>C. perfringens</i> (CFU/100mL)	HF183 (Human) (Copies/100mL)	BacHum (Human) (Copies/100mL)	$\delta^{15}\text{N}$ – $\text{NO}_3^-$ (‰)	$\delta^{18}\text{O}$ – $\text{NO}_3^-$ (‰)
High Elevation Wells	3	15 $\pm$ 9	0 $\pm$ 0	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	4.47 $\pm$ 0.21	1.98 $\pm$ 0.43
Mid Elevation Wells	6	5 $\pm$ 0	0 $\pm$ 0	0.00 $\pm$ 0.00	0.00 $\pm$ 0.00	6.37 $\pm$ 0.40	1.23 $\pm$ 0.32
Low Elevation Wells	2	119 $\pm$ 67	0 $\pm$ 0	0.00 $\pm$ 0.00	0.05 $\pm$ 0.04	6.29 $\pm$ 0.70	2.85 $\pm$ 0.60
Resort Shoreline	4	32 $\pm$ 14	0 $\pm$ 0	n/a	n/a	5.16 $\pm$ 0.31	2.61 $\pm$ 0.70
Puakō Shoreline	16	410 $\pm$ 83	4 $\pm$ 1	0.58 $\pm$ 0.16	0.71 $\pm$ 0.17	8.64 $\pm$ 0.25	7.92 $\pm$ 0.02
Puakō Ponds	2	8165 $\pm$ 2935	18 $\pm$ 11	0.47 $\pm$ 0.28	1.74 $\pm$ 0.55	8.59 $\pm$ 0.41	4.19 $\pm$ 0.53

Number of observations per location, per parameter:

High Elevation Wells: *Enterococcus* spp. = 6, *C. perfringens* = 6, HF183 (Human) = 6, BacHum (Human) = 6,  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  = 9,  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  = 9

Mid Elevation Wells: *Enterococcus* spp. = 16, *C. perfringens* = 16, HF183 (Human) = 11, BacHum (Human) = 11,  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  = 21,  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  = 21


Low Elevation Wells: *Enterococcus* spp. = 6, *C. perfringens* = 6, HF183 (Human) = 4, BacHum (Human) = 4,  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  = 7,  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  = 7

Resort Shoreline: *Enterococcus* spp. = 12, *C. perfringens* = 12,

Puakō Shoreline: *Enterococcus* spp. = 137, *C. perfringens* = 137, HF183 (Human) = 48, BacHum (Human) = 48,  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  = 89,  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  = 89

Puakō Ponds: *Enterococcus* spp. = 7, *C. perfringens* = 5, HF183 (Human) = 4, BacHum (Human) = 4,  $\delta^{15}\text{N}$  –  $\text{NO}_3^-$  = 7,  $\delta^{18}\text{O}$  –  $\text{NO}_3^-$  = 7

mL; range: 324 - 19,018 MPN/100 mL) (Fig. 1A, Table 1). These ponds had values that consistently exceeded the HDOH statistical threshold standard of 130 MPN/100 mL. The second highest *Enterococcus* spp. concentrations were found along Puakō's shoreline ( $410 \pm 83$  MPN/100 mL; 0-7985 MPN/100 mL), with concentrations on average exceeding the HDOH statistical threshold standard. *Enterococcus* spp. concentrations were an order of magnitude lower at the resorts than those measured along the Puakō shoreline and anchialine ponds, and lower than state standards (Table 1). Epidemiological studies have shown that a swimmer's chance of contracting gastroenteritis is 3.6% at *Enterococcus* spp. concentrations of 35 MPN/100 mL (reviewed in Fujioka et al 2015). *Clostridium perfringens* concentrations were similar among high elevation wells ( $0 \pm 0$  CFU/100 mL), mid elevation wells ( $0 \pm 0$  CFU/100 mL), low elevation wells ( $0 \pm 0$  CFU/100 mL), and at the resorts shoreline ( $0 \pm 0$  CFU/100 mL) (Fig. 1B, Table 1). The average *C. perfringens* concentration along the Puakō shoreline ( $4 \pm 1$  CFU/100 mL) was similar to the groundwater wells and resorts shoreline averages. However, the range for *C. perfringens* concentrations along Puakō's shoreline was greater (0 - 90 CFU/100mL) compared to ranges for groundwater wells and resorts' shoreline waters. *C. perfringens* concentrations in Puakō's anchialine ponds ( $18 \pm 11$  CFU/100 mL; 0 - 52 CFU/100 mL) exceeded the recommended standard to HDOH of 5 CFU/ 100 mL for marine recreational waters (Fujioka et al. 1997) and were within the range reported for non-point source sewage pollution (Fung et al. 2007). Human-associated *Bacteroides* (using two different molecular markers HF183 and BacHum) were the highest along Puakō's shoreline and within the Puakō's anchialine ponds (Fig. 1C, Table 1). A recent study found that a swimmer had 1.2 – 3% chance of contracting gastroenteritis at BacHum levels between 1.7 – 3.6 copies/ 100 mL (Boehm et al. 2015) – the levels observed within Puakō's anchialine ponds and shoreline waters (Fig. 1C).




**Figure 2.** Spatial distribution of  $\delta^{15}\text{N}$  macroalgae values and nutrient concentrations along the Puakō shoreline, and at four adjacent resorts north (Mauna Kea, Hapuna Prince) and south (Fairmont Orchid, Mauna Lani) of the community on Hawai'i Island. A)  $\delta^{15}\text{N}$  macroalgae, B) Total Dissolved Nitrogen (TDN), C) Total Dissolved Phosphorus (TDP), and D) dissolved silica ( $\text{H}_4\text{SiO}_4$ ). Samples were collected from June – August 2016, and January to March 2017. Averages ( $\pm$ SE) for parameters can be found in Figure 3 and Table 2.

Seven stations (station #: 2, 3, 4, 5, 6, 7, and 14) along the Puakō shoreline had  $\delta^{15}\text{N}$  macroalgal values  $+7\text{‰}$  or greater, which is considered to be within the range of values measured for sewage (reviewed in Wiegner et al. 2016). At the resorts,  $\delta^{15}\text{N}$  macroalgal values were lower than those

observed at many of the Puakō shoreline stations (Fig. 2A), with values indicative of  $\text{NO}_3^-$  from upper and lower elevational wells, well fertilized soils and ones under Kiawe trees, and ocean water (Fig. 3). However, it is important to note that past studies have found that macroalgae assimilate N more rapidly under low  $\text{NO}_3^-$  concentrations (Fujita 1985), and that  $\delta^{15}\text{N}$  in macroalgal tissues can be underestimated by up to 6‰ in waters with high  $\text{NO}_3^-$  concentrations ( $> 10 \mu\text{mol/L}$ ) (Swart et al. 2014). All shoreline stations had  $\text{NO}_3^- + \text{NO}_2^-$  concentrations that exceeded this value. If the  $\delta^{15}\text{N}$  macroalgal values are adjusted for possible increased N isotope discrimination at higher  $\text{NO}_3^-$  concentrations, then algal shoreline values fall within the range reported for sewage ( $> +7\text{‰}$ , reviewed in Wiegner et al. 2016).

There were no consistent nutrient concentration patterns across sampling locations (Table 2). However, the greatest variability in nutrient concentrations ( $\text{NO}_3^- + \text{NO}_2^-$ , TDN, and  $\text{PO}_4^{3-}$ ) was generally observed along Puakō's shoreline and



**Table 2.** Average  $\pm$  SE of salinity, turbidity (NTU), and nutrient concentrations ( $\mu\text{mol/L}$ ) in water samples for high elevation wells (Waikoloa Village), mid elevation wells (Mauna Lani), low elevation wells (Puakō wells), resort shoreline stations (Mauna Kea, Hapuna Prince, Fairmont Orchid, and Mauna Lani), Puakō shoreline stations (low and high salinity stations), and Puakō ponds (anchialine ponds). Number of stations sampled within each watershed category varied between 3-16 (station n) and the total number of observations per location per parameter are footnoted below.

Watershed Location	Station (n)	Salinity	Turbidity	$\text{NO}_3^- + \text{NO}_2^-$	$\text{NH}_4^+$	TDN	$\text{PO}_4^{3-}$	$\text{H}_4\text{SiO}_4$
High Elevation Well	3	$0.18 \pm 0.01$	$0.39 \pm 0.07$	$95.27 \pm 2.33$	$2.29 \pm 0.89$	$102 \pm 1$	$2.35 \pm 0.06$	$775 \pm 53$
Mid Elevation Well	6	$1.76 \pm 0.09$	$0.84 \pm 0.15$	$120.35 \pm 4.62$	$1.46 \pm 0.36$	$133 \pm 8$	$1.54 \pm 0.05$	$760 \pm 28$
Low Elevation Well	2	$1.91 \pm 0.27$	$6.88 \pm 2.74$	$114.55 \pm 8.93$	$3.31 \pm 1.12$	$127 \pm 15$	$1.94 \pm 0.57$	$657 \pm 76$
Resorts Shoreline	4	$22.37 \pm 2.93$	$1.11 \pm 0.18$	$47.01 \pm 9.68$	$0.80 \pm 0.18$	$57 \pm 11$	$1.01 \pm 0.19$	$328 \pm 77$
Puakō Shoreline	16	$21.53 \pm 0.76$	$0.23 \pm 0.02$	$65.51 \pm 5.21$	$1.64 \pm 0.10$	$77 \pm 5$	$1.60 \pm 0.15$	$447 \pm 88$
Puakō Pond	2	$5.29 \pm 0.61$	$10.62 \pm 2.33$	$110.32 \pm 13.48$	$3.94 \pm 1.24$	$120 \pm 17$	$1.79 \pm 0.50$	$655 \pm 90$

Number of observations per location, per parameter:

High Elevation Wells: Salinity = 8, Turbidity = 6,  $\text{NO}_3^- + \text{NO}_2^-$  = 11,  $\text{NH}_4^+$  = 11, TDN = 6,  $\text{PO}_4^{3-}$  = 11,  $\text{H}_4\text{SiO}_4$  = 8

Mid Elevation Wells: Salinity = 20, Turbidity = 16,  $\text{NO}_3^- + \text{NO}_2^-$  = 25,  $\text{NH}_4^+$  = 25, TDN = 16,  $\text{PO}_4^{3-}$  = 25,  $\text{H}_4\text{SiO}_4$  = 20

Low Elevation Wells: Salinity = 6, Turbidity = 6,  $\text{NO}_3^- + \text{NO}_2^-$  = 8,  $\text{NH}_4^+$  = 8, TDN = 6,  $\text{PO}_4^{3-}$  = 8,  $\text{H}_4\text{SiO}_4$  = 6

Resort Shoreline: Salinity = 12, Turbidity = 12,  $\text{NO}_3^- + \text{NO}_2^-$  = 12,  $\text{NH}_4^+$  = 12, TDN = 12,  $\text{PO}_4^{3-}$  = 12,  $\text{H}_4\text{SiO}_4$  = 12


Puakō Shoreline: Salinity = 137, Turbidity = 137,  $\text{NO}_3^- + \text{NO}_2^-$  = 135,  $\text{NH}_4^+$  = 135, TDN = 135,  $\text{PO}_4^{3-}$  = 135,  $\text{H}_4\text{SiO}_4$  = 135

Puakō Ponds: Salinity = 6, Turbidity = 2,  $\text{NO}_3^- + \text{NO}_2^-$  = 7,  $\text{NH}_4^+$  = 7, TDN = 6,  $\text{PO}_4^{3-}$  = 7,  $\text{H}_4\text{SiO}_4$  = 6

anchialine ponds (Table 2). Nutrient concentrations in waters fronting the resorts were comparable to those observed along the Puakō shoreline (Table 2), except that Puakō had more stations with higher concentrations of TDN and TDP (Fig. 2B, C).  $H_4SiO_4$  concentrations varied along the shoreline, with high concentrations indicative of areas of submarine groundwater discharge (SGD) that were both located within

Puakō and in front of the resorts (Fig. 2D).

To further examine sources of  $NO_3^-$  to mid-elevation groundwater wells, low elevation groundwater wells, anchialine ponds, and shoreline waters, bi-plots of  $\delta^{15}N$ - and  $\delta^{18}O$ - $NO_3^-$  were created, plotting



averages for each station relative to averages for all potential  $NO_3^-$  sources (Fig. 4). This step allowed for the stable isotope data for samples to be visualized relative to values for the sources. These figures were used to decide on which sources to include in the Stable Isotope Analysis in R (SIAR, v. 4.0) mixing models. They also allowed us to determine if different sources have overlapping stable isotope values. This is important because if there is overlap, it is harder to distinguish each source's contribution to a sample from one another in the modeling effort. Based on this analysis, we decided to include sewage, groundwater, and kiawe/fertilized soil. Note that ocean water was not included as a  $NO_3^-$  source as its  $NO_3^-$  concentrations were below detection limits for stable isotope analysis ( $> 2 \mu\text{mol/L}$ , Coplen et al. 2012).


**Table 3.** Mean  $\pm$  SE of  $\delta^{15}N$ - and  $\delta^{18}O$ -  $NO_3^-$  (‰) and nutrient concentrations ( $\mu\text{mol/L}$ ) of potential  $NO_3^-$  sources (high elevational groundwater, sewage, and kiawe/fertilized soil) collected in Puakō as part of this study and a previous one (Abaya et al. 2018). Sources and nutrients without SE indicate single sample (n=1).

Source	$\delta^{15}N$ $NO_3^-$	$\delta^{18}O$ $NO_3^-$	$NO_2^- + NO_3^-$	$NH_4^+$	TDN	$PO_4^{3-}$	$H_4SiO_4$
Groundwater	$4.08 \pm 0.31$	$1.57 \pm 0.44$	$95.27 \pm 2.33$	$2.29 \pm 0.89$	$102 \pm 1$	$2.35 \pm 0.06$	$775 \pm 53$
Sewage	$11.73 \pm 1.16$	$14.62 \pm 4.77$	$49.06 \pm 33.90$	$4803.99 \pm 1160.31$	$2787 \pm 1602$	$307.21 \pm 47.29$	$467 \pm 97$
Soil	$2.57 \pm 1.74$	$1.56 \pm 2.13$	$4775.26 \pm 3051.69$	$446.41 \pm 162.12$	3	$145.33 \pm 111.11$	1

To determine the percent contribution of potential  $NO_3^-$  sources to the various water types examined in this project, we used SIAR. Both  $\delta^{15}N$ - and  $\delta^{18}O$ -  $NO_3^-$  were used in the models, and the potential  $NO_3^-$  sources examined were sewage, groundwater, and kiawe/fertilized soil (Table 3). SIAR was first used to examine potential  $NO_3^-$  sources to mid-elevation groundwater wells (MEW), low-elevation groundwater wells (LEW), Puakō's anchialine ponds (Ponds), the resorts' shoreline waters (Resorts), and Puakō's shoreline waters (Fig. 5, Table 4). Note, in this model, data from all stations within a water type were averaged and station differences within water types were not examined. The

second SIAR model then examined potential  $\text{NO}_3^-$  sources to Puakō's and the resorts' shoreline stations (Fig. 6, Table 5). This analysis was done to identify any shoreline sewage pollution hot spots. The Shapiro-Wilk test for normality was used to verify that isotope data for the three  $\text{NO}_3^-$  sources were normally distributed.  $\delta^{15}\text{N}$ - and  $\delta^{18}\text{O}-\text{NO}_3^-$  data were normally distributed for all sources. Relative percent contributions of each  $\text{NO}_3^-$  source to each water type or station are reported as 50% Bayesian credibility interval. This credibility interval allows a wider range of natural variability and uncertainty to be reported within a system with multiple sources (Parnell et al. 2010).

Sewage contributions to the  $\text{NO}_3^-$  pool varied among water types (Fig. 5, Table 4). Puakō's anchialine ponds had the largest sewage contribution to the  $\text{NO}_3^-$  pool (36 – 55%), followed by Puakō's shoreline waters (23 – 26%), resort shoreline waters (12 – 17%), low elevational groundwater wells (9 – 20%), and mid-elevational groundwater wells (1 – 4%). Along Puakō's shoreline, ten stations (station #: 1, 2, 3, 4, 5, 6, 11, 12, 13, and 14) had sewage contributing up to 40% or more (high end of the range) to the shoreline  $\text{NO}_3^-$  (Fig. 6, Table 5). Several of these stations have been identified as sewage pollution hotspots either from  $\delta^{15}\text{N}$  macroalgal values, sewage pollution scores, or dye tracer tests (Abaya et al. 2018, Wiegner et al. unpubl. data)



**Figure 5.** Average proportional contribution of sewage to the  $\text{NO}_3^-$  pool in different water types measured in Puakō's watershed, Hawai'i. Proportions were estimated using SIAR (v. 4.0). Boxplots illustrate the 50<sup>th</sup>, 75<sup>th</sup>, and 95<sup>th</sup> percentiles from light to dark.


**Table 4.** The range of proportional contributions (%) of potential  $\text{NO}_3^-$  sources (sewage, high groundwater, kiawe/fertilized soil) within five different water types: mid elevation wells (MEW), low elevation wells (LEW), anchialine ponds within Puakō (Ponds), shoreline waters at resorts adjacent to Puakō (Resorts), and shoreline waters at Puakō. Percent contributions are reported as the 50% Bayesian Credibility interval and analyses were run in SIAR (v. 4.0).

Water Type	Sewage	Groundwater	Soil
MEW	1 – 4	84 - 97	0 – 10
LEW	9 - 20	38 - 64	20 - 43
Ponds	36 - 55	31 - 51	1 - 20
Resorts	12 - 17	59 - 81	3 - 24
Puakō Shoreline	23 - 26	66 - 75	0 – 7

**Objective 2:** Quantify water quality impairment caused by homes with septic tanks and aerobic treatment units (ATU) through dye tracer studies, and measurements of macroalgal tissue  $\delta^{15}\text{N}$ , FIB (including *Bacteroides*), and nutrients (including  $\delta^{15}\text{N}$  – and  $\delta^{18}\text{O}-\text{NO}_3^-$ ).

**Accomplishments and ongoing efforts:** In late 2016, we informed homeowners of our interest in doing dye tracer studies through a community email newsletter (Clean Water for Reefs - Puakō newsletter),


an announcement at a community meeting, and through direct interaction with homeowners. Homeowners interested in participating in the dye tracer studies contacted us, and we discussed the experimental process with them. We completed all six dye tracer tests. They were conducted at homes with septic tanks (n = 3) and ATUs (n = 3) (February - May 2017). This included two homes on the landward (mauka) side of the street. Note, dye tracer tests were conducted for cesspools in an earlier project (NOAA CRCP). Analyses of field samples from the dye tracer tests have been completed. Macroalgae and water quality samples were collected in front of all of the dye tracer locations from June – August 2017. FIB,  $\delta^{15}\text{N}$  macroalgae tissue, and nutrient samples have all been analyzed. Analysis of  $\delta^{15}\text{N}$ - and  $\delta^{18}\text{O}$ - of  $\text{NO}_3^-$  in water samples is complete and a SIAR model was



**Table 5.** The range of proportional contributions (%) of potential  $\text{NO}_3^-$  sources (sewage, high elevation groundwater, and kiawe/fertilized soils) at shoreline stations in Puakō (Stations: 1-16) and adjacent resorts (Stations: 17-20) in Hawai'i. Percent contributions are reported as the 50% Bayesian credibility interval. Analysis was conducted in SIAR (v. 4.0)

Location	Station #	Sewage	Groundwater	Soil
Puakō	1	28 - 50	27 - 50	16 - 40
	2	5 - 52	26 - 51	16 - 41
	3	22 - 41	25 - 48	20 - 43
	4	24 - 43	23 - 46	19 - 43
	5	24 - 44	23 - 47	19 - 43
	6	26 - 47	21 - 45	14 - 39
	7	13 - 35	27 - 52	21 - 45
	8	9 - 25	29 - 53	27 - 49
	9	14 - 28	33 - 57	20 - 43
	10	12 - 24	35 - 60	19 - 42
	11	33 - 45	24 - 45	33 - 45
	12	31 - 45	24 - 45	20 - 41
	13	27 - 48	21 - 44	14 - 38
	14	27 - 47	28 - 52	15 - 40
	15	24 - 36	27 - 48	23 - 42
	16	11 - 22	35 - 60	21 - 43
Mauna Kea Beach	17	17 - 29	32 - 55	21 - 42
Hapuna Beach Hotel	18	2 - 12	35 - 58	30 - 51
Fairmont Orchid	19	2 - 16	35 - 63	23 - 47
Mauna Lani Beach	20	17 - 33	33 - 56	20 - 42

utilized to determine the relative percent contributions of  $\text{NO}_3^-$  sources in front of homes with different



OSDS. Analysis of *Bacteroides* samples is still pending.

**Findings:** Dye was observed at the shoreline in five of the six tests. The time it took the dye to travel from the septic tank or ATU to the shoreline varied from 5 hours to 11 days, with flow rates from 3 to 137 m/d. In general, dye appeared at the shoreline sooner in front of homes with drainage fields closer to the shoreline. The type of system (septic tank vs. ATU) did not affect how fast dye reached

**Table 6.** Average ( $\pm$ SE) water quality condition in front of homes with different on-site sewage disposal systems (OSDS: cesspool, septic tanks, aerobic treatment units (ATU)) at Puakō, Hawai‘i. Measurements were made from November 2014 - July 2017. OSDS n = number of OSDS within a type sampled. Samples n = number of samples collected in front of each home.

OSDS type	OSDS n	Sample n	Clostridium		$\delta^{15}\text{N}$ Algae (%)	$\delta^{15}\text{N}$ - $\text{NO}_3^-$ (%)	$\text{NO}_3^- + \text{NO}_2^-$ ( $\mu\text{mol/L}$ )	$\text{NH}_4^+$ ( $\mu\text{mol/L}$ )	TDN ( $\mu\text{mol/L}$ )	$\text{PO}_4^{3-}$ ( $\mu\text{mol/L}$ )	TDP ( $\mu\text{mol/L}$ )	$\text{H}_4\text{SiO}_4$ ( $\mu\text{mol/L}$ )
			<i>Enterococcus</i> spp. (MPN/100 mL)	<i>perfringens</i> (CFU/100 mL)								
Cesspool	4	4 - 10	239 $\pm$ 56	4 $\pm$ 1	9.12 $\pm$ 0.37	11.07 $\pm$ 0.43	105.88 $\pm$ 14.98	2.01 $\pm$ 0.22	116 $\pm$ 15	3.18 $\pm$ 0.43	3.62 $\pm$ 0.47	325 $\pm$ 34
Septic tank	3	3	4182 $\pm$ 3772	7 $\pm$ 3	7.04 $\pm$ 0.45	7.55 $\pm$ 0.41	117.14 $\pm$ 20.22	3.00 $\pm$ 1.28	125 $\pm$ 23	2.62 $\pm$ 0.40	3.09 $\pm$ 0.48	439 $\pm$ 44
ATU	3	3	439 $\pm$ 244	3 $\pm$ 1	8.19 $\pm$ 1.18	9.03 $\pm$ 1.09	157.51 $\pm$ 41.27	3.42 $\pm$ 0.71	165 $\pm$ 45	3.86 $\pm$ 0.93	4.21 $\pm$ 1.05	588 $\pm$ 86

the shoreline, with ATUs having both the slowest and fastest flow rates. These results suggest that the underlying geology (e.g., presence or absence of large fractures in the basalt) likely controls how fast sewage flows from the OSDS to the shoreline.

The only test where dye was not observed at the shoreline was from the home farthest landward (mauka). Here, the drainage field was 122 m from the shoreline; at the other five homes, this averaged 40 m. Additionally, this home was located in the northern part of the neighborhood, where fewer groundwater springs have been observed. It is possible that we either missed the spring where


the dye emerged, or that we did not sample long enough to capture the dye emerging from the spring (after 16 days, flow rate of <8 m/d).

Concentrations of FIB and nutrients were similar in front of homes with different OSDS (Fig. 7). The average concentration of *Enterococcus* spp. in front of homes with differing OSDS all exceeded the HDOH statistical threshold value of 130 MPN/100 mL (Fig. 7A). *Enterococcus* spp. concentrations ranged from 0 – 34,330 MPN/100 mL. The average concentration of *C. perfringens* in front of the homes with differing OSDS encompassed the standard recommended to HDOH for marine


recreational waters of 5 CFU/100 mL (Fujioka et al. 1997), but they were not higher than the range reported for non-point sewage pollution of 10 -100 CFU/100 mL (Fig. 7B, Fung et al. 2007). *C. perfringens* concentrations ranged from 0 – 27 CFU/100 mL.  $\delta^{15}\text{N}$  in macroalgae was similar in front of homes with the different OSDS types (Fig. 8). The average  $\delta^{15}\text{N}$  value in front of homes with cesspools was  $9.12 (\pm 0.37, 4.90 - 12.78)$ , septic tanks  $7.04 (\pm 0.45, 4.84 - 8.70)$ , and ATUs  $8.19 (\pm 1.18, 2.75 - 10.88)$ . Average nutrient concentrations in front of all homes were high. TDN average concentrations in front of homes with different OSDS were all greater than  $100 \mu\text{mol/L}$ , with individual measurements ranging from  $13 - 415 \mu\text{mol/L}$  (Fig 7C).  $\text{NO}_3^- + \text{NO}_2^-$  comprised most of the TDN. TDP averages in front of the homes with differing OSDS were all greater than  $3 \mu\text{mol/L}$ , with individual measurements ranging from  $0.25 - 10.84 \mu\text{mol/L}$  (Fig. 7C).  $\text{PO}_4^{3-}$  comprised most of the TDP.  $\text{H}_4\text{SiO}_4$  concentrations did differ in front of the homes with different OSDS, but this difference was most likely from SGD and not from sewage as  $\text{H}_4\text{SiO}_4$  is an indicator for groundwater (data not shown).

SIAR modeling was used to partition potential  $\text{NO}_3^-$  sources in front of homes with different OSDS types. Details on model effort are described in the results section for Objective 1.

Proportional contributions of sewage to the  $\text{NO}_3^-$  pool in front of homes with different OSDS types varied (Fig. 9). Sewage comprised 25 – 35% of the  $\text{NO}_3^-$  in front of homes with cesspools, 41 – 49% in front of homes with septic tanks, and 12 – 23% in front of homes with ATUs (Table 7).



**Figure 8.** Average ( $\pm$ SE)  $\delta^{15}\text{N}$  macroalgae in front of homes with different onsite sewage disposal systems (OSDS: cesspools, septic tanks, aerobic treatment units (ATU)) in Puakō, Hawai'i. Results from one-way ANOVA are shown on figure.



**Figure 9.** Proportional contributions of sewage as a  $\text{NO}_3^-$  source for three OSDS (cesspool, septic tanks, ATUs) estimated using SIAR (v. 4.0). Boxplots illustrate the 50<sup>th</sup>, 75<sup>th</sup>, and 95<sup>th</sup> percentiles from light to dark.

**Objective 3:** Assess whether the proposed sewage treatment upgrades are sufficient for meeting water quality standards.

**Accomplishments and ongoing efforts:** Data from objectives 1 and 2 have been finalized. A three end-member mixing model was used to determine if sewage treatment upgrades are sufficient for meeting water quality standards for FIB.

**Table 7.** The range of proportional contributions (%) of potential NO<sub>3</sub><sup>-</sup> sources (sewage, high elevation groundwater, and kiawe/fertilized soils) at three different on-site disposal systems (OSDS) in Puakō: cesspools, septic tanks, and ATUs. Percent contributions are reported as the 50% Bayesian credibility interval. Analysis was conducted in SIAR (v. 4.0)

OSDS Type	Sewage	Groundwater	Soil
Cesspools	25 – 35	33 – 53	14 – 35
Septic Tanks	41 – 49	30 – 49	3 – 20
ATUs	12 - 23	36 - 58	23 - 44

**Findings:** A three end-member mixing model (sewage, groundwater, seawater) was used to estimate the fraction of FIB (*Enterococcus* spp. and *C. perfringens*) from sewage in groundwater spring samples along Puakō’s shoreline, and to evaluate potential shoreline water quality improvements that will occur by switching from the current distribution of OSDS to all septic systems, all ATUs, or a sewage treatment plant. FIB concentrations measured in groundwater, sewage, and ocean water from this study and our previous NOAA CRCP grant were used in these calculations. The sewage end-member was assumed to be a mixture of water from different OSDS, with 44% cesspool, 48% septic

**Table 8.** Calculated concentration of fecal indicator bacteria (*Enterococcus* spp. and *Clostridium perfringens*) at shoreline under different scenarios of neighborhood-wide OSDS conversion (see Objective 3 for details).

Station	<i>Enterococcus</i> spp. (MPN/100 mL)				<i>C. perfringens</i> (CFU/100 mL)			
	Today	Septic	ATU*	Treatment*	Today	Septic	ATU	Treatment
1	14	10	10	1.3	1.38	0.72	0.72	0.01
2	97	68	68	3.9	9.72	5.03	5.03	0.00
3	272	190	190	2.7	3.71	1.92	1.92	0.00
4	155	108	108	2.3	3.67	1.90	1.90	0.00
5	645	448	448	0.5	2.14	1.11	1.11	0.01
6	408	283	283	0.5	2.58	1.34	1.34	0.01
7	177	123	123	1.2	5.47	2.83	2.83	0.01
8	426	296	296	2.1	4.10	2.12	2.12	0.00
9	360	251	251	2.0	1.17	0.61	0.61	0.00
10	77	54	54	2.7	2.90	1.50	1.50	0.00
11	485	337	337	0.5	5.11	2.65	2.65	0.01
12	396	275	275	0.8	3.10	1.60	1.60	0.01
13	1931	1341	1341	0.5	1.86	0.96	0.96	0.01
14	248	173	173	3.6	6.81	3.52	3.52	0.00
15	288	200	200	0.1	6.23	3.23	3.23	0.01
16	420	292	292	2.0	1.54	0.80	0.80	0.00

\* ATU = Aerobic Treatment Unit; Treatment = Sewage Treatment Facility

systems and 8% ATUs (Aqua Engineering 2015). FIB concentrations in cesspools were based on measurements of raw sewage at Puakō, which are similar to raw sewage values measured elsewhere (Ahmed et al. 2008, Kay et al. 2008). Septic tanks were assumed to decrease *Enterococcus* spp. and *C. perfringens* concentrations by 50% and 68%, respectively, based on published studies (Chauret et al. 1999, Withers et al. 2011). We found no published information regarding the effectiveness of ATUs in reducing FIB concentrations in their effluent, so we assumed the same reduction as septic systems. This assumption may underestimate the FIB concentration in the ATU effluent since the residence




removal of sewage from the water table with a treatment plant, the resulting shoreline water would only be a mixture of groundwater and seawater.


#### 4. Deliverables and Outcomes (How did this project address critical management needs?)

**Deliverables:** Ten undergraduate students were trained on this project to date. They include six summer interns (Saria Adnan Sultan, Christopher Thompson, Adel Sharif, Carmen Garson-Shumay, Tyler Gerken, and Amy Olson; Fig. 10) supported by the National Science Foundation (NSF) Research Experiences for Undergraduates (REU) program at University of Hawai'i at Hilo [UH Hilo; Pacific Internship

Programs for Exploring Science (PIPES) – <https://hilo.hawaii.edu/uhintern/>). They presented results from their internships at the end-of-the-summer symposium and wrote final reports. Early in 2017 (February – March 2017), one of the summer interns, Saria Adnan Sultan, presented findings from her internship in a poster at the ASLO conference in Honolulu, HI



**Figure 12.** Adel Sharif's (summer 2017 intern) poster presented at the Association for the Sciences of Limnology and Oceanography (ASLO) conference in Victoria, Canada (June 2018). His travel was supported by UH Hilo PIPES program.



(Fig. 11). Her travel was supported by ASLO Minorities Program (ASLOMP) and UH Hilo PIPES program. This past June (2018), Adel Sharif, our most recent summer intern, presented findings from his internship on this project at the ASLO Aquatic Sciences conference in Victoria, Canada (Fig. 12). His travel expenses were paid with funds from the UH Hilo PIPES program (NSF REU). We also had several other undergraduates working on this project, including: Carey Demapan (NSF Experimental Program to Stimulate Competitive Research (EPSCoR) Ike Wai scholars program 2016-2017, 2018

grant-employed student assistant), Melia Takakusagi (NIH SHARP intern), and Julia Stuart (grant-employed student assistant). These latter students assisted with field preparations and laboratory sample processing. Both Carey and Melia presented posters on their experiences working on the project in April 2017 as part of their respective internship programs.

We also produced three community handouts for the Puakō community in collaboration with The Nature Conservancy (TNC; Figs. 13 - 15). Two were for Puakō's Wastewater Symposium in January 2017 and 2018 (Figs. 13 and 15) and the other was for a community association meeting in May 2017 (Fig. 14). At this latter meeting, we gave a joint presentation on results from this project with TNC. In December 2017, Dr. Wiegner presented results from this project at the NOAA West Hawaii




Figure 14. Community handout prepared in collaboration with The Nature Conservancy for the Puakō Community Association meeting in May 2017. Meeting was organized by the Coral Reef Alliance.




Figure 15. Community handout prepared in collaboration with The Nature Conservancy for the Puakō Wastewater Forum held January 2018. Meeting was organized by The Coral Reef Alliance.

Symposium in Kona, HI. In January 2018, Dr. Colbert provided an update on this project to the Puakō community at a Coral Reef Alliance sponsored meeting, along with an updated handout (Fig. 15). In 2018, Drs. Wiegner and Colbert gave four presentations on this project at professional meetings: 1) HDOH Cesspool Update (June 2018), 2) Hawai'i Conservation Conference, Honolulu, HI (July 2018), 3) HDOH Joint Government Water Conference, Hilo, HI (August 2018), and 4) HDOH Joint Government Water Conference, Kailua Kona, HI (August 2018).

Outcomes: Results from ASLO poster were shared with the Puakō Community Association at a meeting in May (2017) to provide them with critical information regarding where sewage pollution is entering their aquifer (Fig. 14). An updated handout was shared with the community in January 2018 (Fig. 15). All parameters concur that sewage pollution is largely entering the watershed within the Puakō community, and not from upslope development or adjacent resorts. These findings highlight the need for OSDS (cesspools, septic tanks, and ATU) to be removed in Puakō, and for improved sewage collection and treatment at a centralized facility. This month (September 2018), a request to HDOH to change Puako's priority status for cesspool removal was submitted by Coral Reef Alliance, which included a letter from UH Hilo with data from this and early projects.

5. Obstacles or Delays: Dye tracer tests were delayed due to challenges in finding volunteer homeowners. A point of contact for arranging dye tracer tests with homeowners was established and all six tests have now been conducted.

Results from Northern Arizona University (NAU) for isotope analyses were delayed due to staff shortages and instrumentation downtime.

*Bacteroides* sample analysis is still pending due to staff shortages in the Nelson lab at UH Mānoa.

6. Future needs: There are no future needs as the project is finished.

7. Photo and caption (optional). Photos and captions are contained within figures.

8. Additional Sources of Funding: UH Hilo Seed Funds (from UH Hilo Research Council, Awarded to T. Wiegner), UHH Marine Science Department, UHH Analytical Laboratory (<https://hilo.hawaii.edu/~analab/>), and several UHH internship programs: <sup>a</sup>PIPES (<https://hilo.hawaii.edu/uhintern/>), <sup>b</sup>SHARP (<http://www.uhhilo-sharp.org/>), and <sup>c</sup>NSF EPSCoR Ike Wai (<https://www.hawaii.edu/epscor/>).

9. Field and Laboratory Assistance Provided By:

UH Hilo Technicians: Leilani Abaya, Jazmine Panelo, Caree Edens, and Erin Stamper.

Graduate students: Kristina Remple (UH Mānoa, SOEST).

UH Hilo Undergraduate Interns and Assistants: Saria Adan-Sultan<sup>a</sup>, Christopher Thompson<sup>a</sup>, Adel Sharif<sup>a</sup>, Melia Takakusugi<sup>b</sup>, Carey Demapan<sup>c,d</sup>, Julia Stuart<sup>d</sup>, Bryan Tonga<sup>e</sup>, Carmen Garson-Shumay<sup>a</sup>, Tyler Gerken<sup>a</sup>, and Amy Oleson<sup>a</sup>.

<sup>a</sup> UH Hilo NSF REU PIPES summer interns

<sup>b</sup> UH Hilo NIH SHARP interns

<sup>c</sup> UH Hilo NSF EPSCoR Ike Wai scholars

<sup>d</sup> Student assistant on DAR CRWG project

<sup>e</sup> UH Hilo Analytical Laboratory student assistant

Progress Report Prepared by: Tracy Wiegner, Leilani Abaya, and Steve Colbert

Signature of Point of Contact:



## References:

- Abaya, L.M., T.N. Wiegner, S. Colbert, J. Beets, K.M. Carlson, K.L. Kramer, R. Most, and C. Couch. (2018). A multi-indicator approach for identifying shoreline sewage pollution hotspots adjacent to coral reefs. *Marine Pollution Bulletin* 129(1): 70-80.
- Ahmed, W., Stewart, J., Powell, D. and Gardner, T., (2008). Evaluation of the host-specificity and prevalence of enterococci surface protein (esp) marker in sewage and its application for sourcing human fecal pollution. *Journal of Environmental Quality* 37(4): 1583-1588.
- Aqua Engineering. 2015. Puakō, Hawai'i, Community feasibility study and preliminary engineering report. Prepared for the Coral Reef Alliance.
- Boehm, A.B., Soller, J.A. and Shanks, O.C. (2015). Human-associated fecal quantitative polymerase chain reaction measurements and simulated risk of gastrointestinal illness in recreational waters contaminated with raw sewage. *Environmental Science & Technology Letters* 2(10): 270-275.
- Chauret, C., Springthorpe, S. and Sattar, S., (1999). Fate of *Cryptosporidium* oocysts, *Giardia* cysts, and microbial indicators during wastewater treatment and anaerobic sludge digestion. *Canadian Journal of Microbiology* 45(3): 257-262.
- Coplen, T.B., Haiping, Q., Kinga, R., Casciotti, K., Hannon, J.E. (2012). Determination of the  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  of nitrate in water; RSIL lab code 2900, chap. 17 of Stable isotope-ratio methods, sec. C of Revesz, Kinga, and Coplen, T.B. eds, *Methods of the Reston Stable Isotope Laboratory*; U.S. Geologic Survey Techniques and Methods, book 10, 35 p.
- Fujioka, R.S., B. Roll, and M. Byappanahalli. (1997). Appropriate recreational water quality standards for Hawai'i and other tropical regions based on concentrations of *Clostridium perfringens*. *Proceedings of the Water Environment Federation* 4: 405-411.
- Fujioka, R., Solo-Gabriele, H., Byappanahalli, M. and Kirs, M. (2015). US recreational water quality criteria: a vision for the future. *International Journal of Environmental Research and Public Health* 12(7): 7752-7776.
- Fujita, R.M. (1985) The role of nitrogen status in regulating transient ammonium uptake and nitrogen storage by macroalgae. *Journal of Experimental Marine Biology and Ecology* 92:283-301
- Fung, D.Y.C., R. Fujioka, K. Vijayavel, D. Sato, and D. Bishop. (2007). Evaluation of Fung Double Tube Test for *Clostridium perfringens* and Easyphage test for F-specific RNA coliphages as rapid screening tests for fecal contamination in recreational waters of Hawaii. *Journal of Rapid Methods & Automation in Microbiology* 15: 217-229.
- Kay, D., Crowther, J., Stapleton, C.M., Wyer, M.D., Fewtrell, L., Edwards, A., Francis, C.A., McDonald, A.T., Watkins, J. and Wilkinson, J. (2008). Faecal indicator organism concentrations in sewage and treated effluents. *Water Research* 42(1-2): 442-454.
- Parnell, A., Inger, C.R., Bearhop, S., and Jackson, A.L. (2010) Source partitioning using stable isotopes: coping with too much variation. *PLoS ONE* 5, e9672.
- Swart, P.K., Evans, S., Capo, T., and Altabet, M.A. (2014) The fractionation of nitrogen and oxygen isotopes in macroalgae during the assimilation of nitrate. *Biogeosciences* 11:6147–6157.
- Wiegner, T.N., Mokiao-Lee, A.U., and Johnson, E.E. (2016). Identifying nitrogen sources to thermal tide pools in Kapoho, Hawai'i, U.S.A, using a multi-stable isotope approach. *Marine Pollution Bulletin* 103:63–71.

Withers, P.J.A., Jarvie, H.P. and Stoate, C. (2011). Quantifying the impact of septic tank systems on eutrophication risk in rural headwaters. *Environment International* 37(3): 644-653.