

# ENSO: Recent Evolution, Current Status and Predictions



Update prepared by:  
Climate Prediction Center / NCEP  
2 April 2018

# Outline

Summary

Recent Evolution and Current Conditions

Oceanic Niño Index (ONI)

Pacific SST Outlook

U.S. Seasonal Precipitation and Temperature Outlooks

Summary

# Summary

## ENSO Alert System Status: La Niña Advisory

La Niña conditions are present.\*

Equatorial sea surface temperatures (SSTs) are below average across the central and eastern Pacific Ocean.

A transition from La Niña to ENSO-neutral is most likely (~55% chance) during the March-May season, with neutral conditions likely to continue into the second half of the year.


\* Note: These statements are updated once a month (2<sup>nd</sup> Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).

# Recent Evolution of Equatorial Pacific SST Departures (°C)

From mid April to July 2017, near-to-above average SSTs spanned most of the equatorial Pacific.

During August 2017, above-average SSTs dissipated east of the date line.



Since September 2017, negative SST anomalies have persisted in the east-central equatorial Pacific.



# Niño Region SST Departures (°C) Recent Evolution


The latest weekly SST departures are:

Niño 4	-0.1°C
Niño 3.4	-0.7°C
Niño 3	-0.6°C
Niño 1+2	-0.6°C




# SST Departures ( $^{\circ}\text{C}$ ) in the Tropical Pacific During the Last Four Weeks

During the last four weeks, equatorial SSTs were below average across the central and eastern Pacific Ocean, and above average in the western Pacific.



# Global SST Departures (°C) During the Last Four Weeks


During the last four weeks, equatorial SSTs were above average in the western Pacific. SSTs were below average in the central and eastern Pacific Ocean.



# Weekly SST Departures during the Last Four Weeks

During the last four weeks, below-average SSTs persisted across the east-central equatorial Pacific Ocean. Negative SST anomalies strengthened near the coast of South America.


Weekly SST Anomalies (DEG C)





# Change in Weekly SST Departures over the Last Four Weeks

During the last four weeks, negative changes were observed near the South American coast.






# Upper-Ocean Conditions in the Equatorial Pacific

The basin-wide equatorial upper ocean (0-300 m) heat content is greatest prior to and during the early stages of a Pacific warm (El Niño) episode (compare top 2 panels), and least prior to and during the early stages of a cold (La Niña) episode.

The slope of the oceanic thermocline is least (greatest) during warm (cold) episodes.


Recent values of the upper-ocean heat anomalies (below average) and thermocline slope index (above average) reflect La Niña conditions.

*The monthly thermocline slope index represents the difference in anomalous depth of the 20°C isotherm between the western Pacific (160°E-150°W) and the eastern Pacific (90°-140°W).*




# Central and Eastern Pacific Upper-Ocean (0-300 m) Weekly Average Temperature Anomalies

Positive subsurface temperature anomalies with large fluctuations in amplitude were present from mid-January through mid-July 2017. Negative anomalies lasted from August 2017 to February 2018. Since the end of February, temperature anomalies have been positive and increasing.




# Sub-Surface Temperature Departures in the Equatorial Pacific

In the last two months, positive subsurface temperature anomalies have shifted eastward into the east-central Pacific Ocean.



Recently, positive temperature anomalies at depth have shifted eastward to  $\sim 120^\circ\text{W}$ . Negative anomalies persist closer to the surface between  $\sim 170^\circ\text{W}$  and  $80^\circ\text{W}$ .




# Tropical OLR and Wind Anomalies During the Last 30 Days

Positive OLR anomalies (reduced convection and precipitation) were present near the Date Line. Negative OLR anomalies (enhanced convection and precipitation) were apparent over the far western equatorial Pacific Ocean and parts of Indonesia.

Low-level (850-hPa) winds were anomalous westerly over the far western tropical Pacific Ocean. The winds were anomalous easterly over the central and east-central tropical Pacific.

Upper-level (200-hPa) winds were anomalous westerly over most of the tropical Pacific.



# Intraseasonal Variability

Intraseasonal variability in the atmosphere (wind and pressure), which is often related to the Madden-Julian Oscillation (MJO), can significantly impact surface and subsurface conditions across the Pacific Ocean.

Related to this activity:

Significant weakening of the low-level easterly winds usually initiates an eastward-propagating oceanic Kelvin wave.

# Weekly Heat Content Evolution in the Equatorial Pacific


From August 2017- early January 2018, negative subsurface anomalies persisted in the central and eastern Pacific Ocean.

From December 2017- February 2018, a downwelling Kelvin wave contributed to the eastward shift of above-average subsurface temperatures.

From mid January 2018 to present, an upwelling Kelvin wave resulted in below-average subsurface temperatures in the central and east-central Pacific.

Since early February 2018, another downwelling Kelvin wave has led to positive subsurface anomalies as far east as  $\sim 120^{\circ}\text{W}$ .

Equatorial oceanic Kelvin waves have alternating warm and cold phases. The warm phase is indicated by dashed lines. Downwelling and warming occur in the leading portion of a Kelvin wave, and up-welling and cooling occur in the trailing portion.



# Low-level (850-hPa) Zonal (east-west) Wind Anomalies ( $\text{m s}^{-1}$ )


Through mid-January 2018, easterly wind anomalies prevailed across the central equatorial Pacific.

Up to late February 2018, the Madden Julian Oscillation (MJO) contributed to the eastward propagation of low-level wind anomalies.

Since early March 2018, easterly (westerly) wind anomalies have been more persistent over the east-central (western) Pacific Ocean.

Westerly Wind Anomalies (orange/red shading)

Easterly Wind Anomalies (blue shading)






# Upper-level (200-hPa) Velocity Potential Anomalies

Until April 2018, anomalous upper-level divergence (green shading) generally persisted near Indonesia.

Eastward propagation of regions of upper-level divergence (green shading) and convergence (brown shading) have been evident through early March 2018.

Unfavorable for precipitation (brown shading)  
Favorable for precipitation (green shading)


Note: Eastward propagation is not necessarily indicative of the Madden-Julian Oscillation (MJO).



# Outgoing Longwave Radiation (OLR) Anomalies

At least since October 2017, positive OLR anomalies generally persisted over the central Pacific Ocean. At the same time, negative OLR anomalies were intermittent near the Maritime Continent.

Drier-than-average Conditions (orange/red shading)  
Wetter-than-average Conditions (blue shading)



# Oceanic Niño Index (ONI)

The ONI is based on SST departures from average in the Niño 3.4 region, and is a principal measure for monitoring, assessing, and predicting ENSO.

Defined as the three-month running-mean SST departures in the Niño 3.4 region. Departures are based on a set of improved homogeneous historical SST analyses (Extended Reconstructed SST - ERSST.v5). The SST reconstruction methodology is described in Huang et al., 2017, J. Climate, vol. 30, 8179-8205.)

It is one index that helps to place current events into a historical perspective

# NOAA Operational Definitions for El Niño and La Niña

El Niño: characterized by a positive ONI greater than or equal to  $+0.5^{\circ}\text{C}$ .

La Niña: characterized by a negative ONI less than or equal to  $-0.5^{\circ}\text{C}$ .


By historical standards, to be classified as a full-fledged El Niño or La Niña episode, these thresholds must be exceeded for a period of at least 5 consecutive overlapping 3-month seasons.

CPC considers El Niño or La Niña conditions to occur when the monthly Niño3.4 OISST departures meet or exceed  $\pm 0.5^{\circ}\text{C}$  along with consistent atmospheric features. These anomalies must also be forecasted to persist for 3 consecutive months.

# ONI (°C): Evolution since 1950

The most recent ONI value (December 2017- February 2018) is  $-0.9^{\circ}\text{C}$ .

El Niño ↑  
Neutral  
La Niña ↓






# CPC/IRI Probabilistic ENSO Outlook

Updated: 8 March 2018

A transition from La Niña to ENSO-neutral is expected during the Northern Hemisphere spring (~55% chance of ENSO-neutral during March-May). Thereafter, ENSO-neutral conditions are likely into the second half of the year.



# IRI/CPC Pacific Niño

## 3.4 SST Model Outlook

The majority of models predict the return to ENSO-neutral by summer 2018, continuing through fall 2018.




Figure provided by the International Research Institute (IRI) for Climate and Society (updated 19 March 2018).




# SST Outlook: NCEP CFS.v2 Forecast (PDF corrected)

Issued: 1 April 2018


The CFS.v2 ensemble mean (black dashed line) favors ENSO-neutral through Northern Hemisphere fall 2018.

CFSv2 forecast Nino3.4 SST anomalies (K) (PDF corrected)




— Latest 8 forecast members  
— Earliest 8 forecast members  
— Other forecast members  
- - - Forecast ensemble mean  
— NCDc daily analysis

(Model bias correct base period: 1999–2010; Climatology base period: 1982–2010)




# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

Since mid February 2018, anomalous ridging over the North Pacific Ocean has been accompanied by an anomalous trough (and below-average temperatures) over the western contiguous United States. Over the eastern United States, the pattern has been variable.




# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days

Since mid February 2018, anomalous ridging over the North Pacific Ocean has been accompanied by an anomalous trough (and below-average temperatures) over the western contiguous United States. Over the eastern United States, the pattern has been variable.



# Atmospheric anomalies over the North Pacific and North America During the Last 60 Days


Since mid February 2018, anomalous ridging over the North Pacific Ocean has been accompanied by an anomalous trough (and below-average temperatures) over the western contiguous United States. Over the eastern United States, the pattern has been variable.




# U.S. Temperature and Precipitation Departures During the Last 30 Days

End Date: 31 March 2018

### Percent of Average Precipitation




### Temperature Departures (degree C)




# U.S. Temperature and Precipitation Departures During the Last 90 Days

End Date: 31 March 2018

Percent of Average Precipitation




Temperature Departures (degree C)



# U. S. Seasonal Outlooks

April - June 2018

The seasonal outlooks combine the effects of long-term trends, soil moisture, and, when appropriate, ENSO.



# Summary

## ENSO Alert System Status: La Niña Advisory

La Niña conditions are present.\*

Equatorial sea surface temperatures (SSTs) are below average across the central and eastern Pacific Ocean.

A transition from La Niña to ENSO-neutral is most likely (~55% chance) during the March-May season, with neutral conditions likely to continue into the second half of the year.

\* Note: These statements are updated once a month (2<sup>nd</sup> Thursday of each month) in association with the ENSO Diagnostics Discussion, which can be found by clicking [here](#).